

**EGM 3401**

**Theory Assignment #2**

**Spring 2017**

Due Date: 15 March 2017

## Question 1

Consider the simple pendulum problem in Chapter 3 on pages 157–163 of the textbook. Using the definition of angular momentum,  ${}^{\mathcal{N}}\mathbf{H}_Q$ , relative to an arbitrary point  $Q$ , show that the component of the reaction force exerted by the hinge in the direction transverse to the arm must be zero.

**Hint:** in order to solve this problem, take the system as being the arm and the particle, then choose appropriate reference point about which to compute the angular momentum.

## Question 2

Let  $\mathcal{A}$  be an inertial reference frame, and let  $\mathcal{B}$  be a reference frame that translates with constant velocity relative to reference frame  $\mathcal{A}$ . Prove that  $\mathcal{B}$  is also inertial reference frame.

## Question 3

Let  $T$  be the kinetic energy of a particle of mass  $m$  and let  $\mathbf{r}$  be the position of the particle measured relative to an inertially fixed point. Suppose now that the position of the particle is parameterized in terms of three independent scalar quantities  $(q_1, q_2, q_3)$  and time, that is,  $\mathbf{r} = \mathbf{r}(q_1, q_2, q_3, t)$ . Finally, let  $\mathbf{v} = \dot{\mathbf{r}} = d\mathbf{r}/dt$  be the inertial velocity and let  $\mathbf{a} = \dot{\mathbf{v}} = d\mathbf{v}/dt = \ddot{\mathbf{r}} = d^2\mathbf{r}/dt^2$  be the inertial acceleration of the particle (that is, assume for this problem that the notation  $d/dt$  refers to a rate of change taken in the inertial reference frame). Prove that

$$\frac{d}{dt} \frac{\partial T}{\partial \dot{q}_j} - \frac{\partial T}{\partial q_j} = \mathbf{F} \cdot \frac{\partial \mathbf{r}}{\partial q_j}, \quad (j = 1, 2, 3),$$

where  $\mathbf{F}$  is the resultant force acting on the particle.

**Hint:** In your solution, use the fact that the inertial velocity can be written as

$$\mathbf{v} = \dot{\mathbf{r}} = \frac{d\mathbf{r}}{dt} = \sum_{i=1}^3 \frac{\partial \mathbf{r}}{\partial q_i} \dot{q}_i + \frac{\partial \mathbf{r}}{\partial t}$$

## Question 4

The moment of momentum of a particle relative to an arbitrary point  $Q$  in an inertial reference frame  $\mathcal{N}$  is defined as

$${}^{\mathcal{N}}\mathbf{L}_Q = (\mathbf{r} - \mathbf{r}_Q) \times m {}^{\mathcal{N}}\mathbf{v},$$

where  $\mathbf{r}$  is the position of the particle measured relative to a point fixed in  $\mathcal{N}$  and  ${}^{\mathcal{N}}\mathbf{v}$  is the velocity of the particle as viewed by an observer fixed in  $\mathcal{N}$ . Show that relative to  $Q$  as

$$\frac{{}^{\mathcal{N}}d}{dt} ({}^{\mathcal{N}}\mathbf{L}_Q) = (\mathbf{r} - \mathbf{r}_Q) \times \mathbf{F} - {}^{\mathcal{N}}\mathbf{v}_Q \times m {}^{\mathcal{N}}\mathbf{v},$$

where  $\mathbf{F}$  is the resultant force acting on the particle.

## Question 5

Consider a system of  $n$  particles of mass  $(m_1, \dots, m_n)$ . Let  $(\mathbf{r}_i, {}^{\mathcal{N}}\mathbf{v}_i, {}^{\mathcal{N}}\mathbf{a}_i)$  be the position, inertial velocity, and inertial acceleration of particle  $i \in [1, \dots, n]$ . Prove the following statements:

(a)  $\sum_{i=1}^n m_i \mathbf{r}_i - m \bar{\mathbf{r}} = \mathbf{0}$ ,

$$(b) \sum_{i=1}^n m_i {}^{\mathcal{N}}\mathbf{v}_i - m {}^{\mathcal{N}}\bar{\mathbf{v}} = \mathbf{0},$$

$$(c) \sum_{i=1}^n m_i {}^{\mathcal{N}}\mathbf{a}_i - m {}^{\mathcal{N}}\bar{\mathbf{a}} = \mathbf{0},$$

where  $m = \sum_{i=1}^n m_i$ , and  $(\bar{\mathbf{r}}, {}^{\mathcal{N}}\bar{\mathbf{v}}, {}^{\mathcal{N}}\bar{\mathbf{a}})$ , are, respectively, the position, velocity, and acceleration of the center of mass of the system.

## Question 6

Consider a system of  $n$  particles of mass  $(m_1, \dots, m_n)$ . Furthermore, let  ${}^{\mathcal{N}}\mathbf{a}_i$ , ( $i = 1, \dots, n$ ) be the acceleration of particle  $i \in [1, \dots, n]$  as viewed by an observer in the inertial reference frame  $\mathcal{N}$ . Finally, let  $\mathbf{R}_i = \mathbf{F}_i + \sum_{j=1}^n \mathbf{f}_{ij}$  be the resultant force acting on particle  $i \in [1, \dots, n]$ , where  $\mathbf{F}_i$  is the resultant *external* force acting on particle  $i$  and  $\mathbf{f}_{ij}$  is the force exerted by particle  $j$  on particle  $i$ . Show that

$$\mathbf{F} = \sum_{i=1}^n \mathbf{F}_i = m {}^{\mathcal{N}}\bar{\mathbf{a}}, \quad (1)$$

where  ${}^{\mathcal{N}}\bar{\mathbf{a}}$  is the acceleration of the center of mass of the system as viewed by an observer in the inertial reference frame  $\mathcal{N}$ .

## Question 7

Consider a system of  $n$  particles of mass  $(m_1, \dots, m_n)$ . Let  $(\mathbf{r}_i, {}^{\mathcal{N}}\mathbf{v}_i, {}^{\mathcal{N}}\mathbf{a}_i)$  be the position, inertial velocity, and inertial acceleration of particle  $i \in [1, \dots, n]$ . Furthermore, let  ${}^{\mathcal{N}}\mathbf{H}_Q$  be the angular momentum of the system relative to an arbitrary point  $Q$ , where  ${}^{\mathcal{N}}\mathbf{H}_Q$  is defined as

$${}^{\mathcal{N}}\mathbf{H}_Q = \sum_{i=1}^n (\mathbf{r}_i - \mathbf{r}_Q) \times m_i ({}^{\mathcal{N}}\mathbf{v}_i - {}^{\mathcal{N}}\mathbf{v}_Q).$$

Show that

$${}^{\mathcal{N}}\mathbf{H}_Q = {}^{\mathcal{N}}\bar{\mathbf{H}} + (\mathbf{r}_Q - \bar{\mathbf{r}}) \times m ({}^{\mathcal{N}}\mathbf{v}_Q - {}^{\mathcal{N}}\bar{\mathbf{v}}),$$

where  $m = \sum_{i=1}^n m_i$  is the total mass of the system and  ${}^{\mathcal{N}}\bar{\mathbf{H}}$  is the angular momentum relative to the center of mass of the system.

## Question 8

Consider a system of  $n$  particles of mass  $(m_1, \dots, m_n)$ . Furthermore, let  ${}^{\mathcal{N}}\mathbf{v}_i$ , ( $i = 1, \dots, n$ ) be the velocity of particle  $i \in [1, \dots, n]$  as viewed by an observer in the inertial reference frame  $\mathcal{N}$ . The angular momentum of a system of  $n$  particles relative to an arbitrary point is defined as

$${}^{\mathcal{N}}\mathbf{H}_Q = \sum_{i=1}^n (\mathbf{r}_i - \mathbf{r}_Q) \times m_i ({}^{\mathcal{N}}\mathbf{v}_i - {}^{\mathcal{N}}\mathbf{v}_Q).$$

Show the following results:

(i)

$$\frac{{}^{\mathcal{N}}d}{dt} ({}^{\mathcal{N}}\mathbf{H}_Q) = \mathbf{M}_Q - (\bar{\mathbf{r}} - \mathbf{r}_Q) \times m {}^{\mathcal{N}}\mathbf{a}_Q,$$

(ii)

$$\frac{{}^{\mathcal{N}}d}{dt} ({}^{\mathcal{N}}\bar{\mathbf{H}}) = \bar{\mathbf{M}},$$

(iii)

$$\frac{{}^{\mathcal{N}}d}{dt} ({}^{\mathcal{N}}\mathbf{H}_O) = \mathbf{M}_O,$$

where  $\mathbf{M}_Q$ ,  $\bar{\mathbf{M}}$ , and  $\mathbf{M}_O$  are, respectively, the moments due to all *external* forces relative to the arbitrary point  $Q$ , the center of mass of the system, and the inertially fixed point  $O$ , and  ${}^{\mathcal{N}}\mathbf{a}_Q$  is the acceleration of point  $Q$  as viewed by an observer in the inertial reference frame  $\mathcal{N}$ .

## Question 10

Consider a system of  $n$  particles of mass  $(m_1, \dots, m_n)$ . Furthermore, let  ${}^{\mathcal{N}}\mathbf{v}_i$ , ( $i = 1, \dots, n$ ) be the velocity of particle  $i \in [1, \dots, n]$  as viewed by an observer in the inertial reference frame  $\mathcal{N}$ . Show that the kinetic energy of the system can be written as

$$T = \frac{1}{2} m {}^{\mathcal{N}}\bar{\mathbf{v}} \cdot {}^{\mathcal{N}}\bar{\mathbf{v}} + \frac{1}{2} \sum_{i=1}^n m_i ({}^{\mathcal{N}}\mathbf{v}_i - {}^{\mathcal{N}}\bar{\mathbf{v}}) \cdot ({}^{\mathcal{N}}\mathbf{v}_i - {}^{\mathcal{N}}\bar{\mathbf{v}}),$$

where  ${}^{\mathcal{N}}\bar{\mathbf{v}}$  is the inertial velocity of the center of mass of the system.